

Please ensure that your abstract fits into one column on one page and complies with the *Instructions to Authors* available from the Abstract Submission web page.

Geochemical, Isotopic and Detrital-Zircon Provenance of Mesozoic Rocks, Oregon and Idaho, U.S.A.

T. A. LAMASKIN^{1*}, J. D. VERVOORT², AND R. J. DORSEY¹

¹1272 University of Oregon, Eugene, OR 97405

(*correspondence: tlamaski@uoregon.edu)

²Wash. State Univ., Webster Physical Science Building, Room 1228, Pullman, WA 99164-6376

The Mesozoic evolution of accreted island arcs and subduction complexes in the Blue Mountain Province (BMP) of eastern Oregon and western Idaho has been explained with a wide variety of tectonic models. Common elements include the intra-oceanic origin for both the Wallowa and Olds Ferry arcs and accretion of the BMP to western Laurentia during Late Jurassic or Middle Cretaceous time.

In this study, we use geochemical, isotopic and detrital-zircon provenance of mudstone-sandstone turbidite couplets from the Blue Mountains to characterize the tectonic settings of sediment source areas and to constrain the timing of terrane accretion. Flat REE patterns, low incompatible element concentrations, positive ϵ_{Nd} values and a lack of Precambrian detrital-zircon grains indicates that during Middle to Late Triassic time, the Wallowa terrane was an intra-oceanic arc system. Conversely, steeper REE patterns, higher incompatible element concentrations, more negative ϵ_{Nd} values and ubiquitous Precambrian detrital-zircon grains indicate a continent-fringing setting for the Triassic Olds Ferry arc. Middle to Late Jurassic deep-marine shales overlying both the Wallowa and Olds Ferry arcs have steepened REE patterns, higher incompatible element concentrations, more negative ϵ_{Nd} values and Precambrian detrital-zircon grains, indicating that following Triassic amalgamation, the entire BMP region received sediment input from continental sources. These results suggest that BMP accretion likely initiated in the Middle Jurassic, earlier than is generally agreed upon.

Our results highlight the need for evaluation of both igneous and coeval sedimentary rocks to adequately characterize ancient tectonic settings. Previous trace-element and isotopic investigations of igneous rocks in the Blue Mountains have suggested an intra-oceanic setting for both the Wallowa and Olds Ferry arcs. Evaluation of contemporaneous sedimentary rocks enhances our understanding of the region and has important implications for Mesozoic tectonic models of western North America.